

STEENHOF, K. 1987. Assessing raptor reproductive success and productivity. Pages 157–170 in B.A. Giron Pendleton, B.A. Millsap, K.W. Cline, and D.M. Bird [Eds.], Raptor management techniques manual. Washington, DC U.S.A.

WARNKE, D.K., D.E. ANDERSEN, C.R. DYKSTRA, M.W. MEY-

ER, AND W.H. KARASOV. 2002. Provisioning rates and time budgets of adult and nestling Bald Eagles at inland Wisconsin nests. *J. Raptor Res.* 36:121–127.

Received 28 April 2001; accepted 31 December 2001

J. Raptor Res. 36(2):139–141

© 2002 The Raptor Research Foundation, Inc.

ABSENCE OF BLOOD PARASITES IN NESTLINGS OF THE ELEONORA'S FALCON (*FALCO ELEONORAE*)

ALEJANDRO MARTÍNEZ-ABRAÍN¹

*Instituto Mediterráneo de Estudios Avanzados IMEDEA (CSIC-UIB), C/Miquel Marqués 21,
07190 Esporles, Mallorca, Spain*

GERARDO URIOS

*Departamento de Biología Vegetal, Facultad de Biología, Universitat de València, Dr. Moliner,
50 Burjassot, Valencia, Spain*

KEY WORDS: *Eleonora's Falcon*; *Falco eleonorae*; *nestlings*; *blood parasites*; *Columbretes Islands*.

Parasites are an important factor influencing the dynamics of populations and the structure of animal communities (Sheldon and Verhulst 1996). In birds, haematozoan parasites have been found in more than 2500 of 4000 species examined (Bennett et al. 1992, Bishop and Bennett 1992). The order Falconiformes includes ca. 285 species (Peirce et al. 1990). No haemoproteids have been described for the families Cathartidae (75 species), Pandionidae (15 species), and Sagittaridae (15 species), but four species of haemoproteids have been described from the family Falconidae (59 species). The Eleonora's Falcon (*Falco eleonorae*) is a migratory falcon that nests on islands of the Mediterranean region and winters in south-east Africa, mainly in Madagascar and the Mascarene islands (Walter 1979).

In this study, we examined blood smears of 42 nestlings of the Eleonora's Falcon (18 in 1999 and 24 in 2000) to detect the presence of blood parasites. The only published works on the prevalence of blood parasites in Eleonora's Falcon are those by Wink et al. (1979) and Ristow and Wink (1985), who reported a low prevalence (13%) of *Leucocytozoon toddi* in adult birds (2 of 16 birds infected), but no information for nestlings was provided. To our knowledge, our work is the first to report on blood parasites in nestlings of the Eleonora's Falcon.

Nestlings sampled came from the Columbretes archipelago, a small (19 ha) volcanic outcrop located 63 km off the coast of Castellón (39°54'N, 0°41'E) where about 30 pairs of Eleonora's Falcon breed (A. Martínez-Abraín unpubl. data). Vegetation is typical of a Mediterranean island with small shrubs and annual plants. The only sources of fresh water are two cisterns which collect water from the scarce rainfall (annual mean ca. 250 mm). All 1999 samples came from the main group of islands (Columbrete Gran and Mancolibre), but in 2000 we included samples from Foradada and Ferrera islands. Nestlings sampled came from eight different nests in 1999 and from 16 in 2000. Blood samples were collected by venipuncture of the ulnar vein of 20–25-d-old chicks from 17–22 September 1999 and from 21–23 September 2000. Smears were air-dried and fixed in methanol on the day of sampling. In the laboratory, slides were stained with Giemsa and examined under a microscope with oil at 1000×, using the techniques of Korpimäki et al. (1995). Prevalence was established through the inspection of 100 fields, containing about 100 erythrocytes each. All smears were inspected twice by the same person (A. Martínez-Abraín) and once by a second observer (B. Esparza) at lower power (400×). We sampled haematophagous night-dwelling insects in September 2000, with a Center for Disease Control (Kimsey and Chaniotis 1984) mosquito trap placed for three consecutive nights on the main island but no haematophagous insect was trapped.

No blood parasites were found in the 42 blood samples taken from Eleonora's Falcon nestlings. Because the Eleonora's Falcon is an island species and marine habitats seem to represent an unsuitable environment for po-

¹ Present address: CPEMN Conselleria de Medio Ambiente, Avda. de los Pinares, 106, 46012 El Saler, Valencia, Spain; E-mail address: jandrom@navegalia.com

tential vectors of haematozoan parasites (Little and Earlé 1994, Piersma 1997) it is very likely that the absence of blood parasites in the nestlings sampled is due to the absence of appropriate vectors in the Columbretes Islands, located far from the mainland. The lack of appropriate vectors is one of the reasons commonly used to explain the absence of blood parasites in birds linked to saline or marine habitats (Greiner et al. 1975, Figuerola et al. 1996) and in Spain birds associated with cliffs, like the Griffon Vulture (*Gyps fulvus*) (Blanco et al. 1998). Tella et al. (1996) attributes the low prevalence of *H. tinnunculi* in adults of the Lesser Kestrel (*Falco naumanni*), breeding in open, arid areas of the northern Iberian plateau, to the scarcity of suitable vectors. In contrast, adult falconids breeding in forested areas may have higher prevalences of infection, as is the case of the American Kestrel (*Falco sparverius*), with a prevalence of 85% (Apanius 1991), 74% for females and 53% for males (Wiehn et al. 1997), and 75% (Castellucci et al. 1998). However, neither Korpimäki et al. (1995) studying Eurasian Kestrels (*Falco tinnunculus*) or Apanius and Kirkpatrick (1988) studying American Kestrels, were able to detect blood parasites in most nestlings, although 69% of juvenile American Kestrels were infected with *H. tinnunculi* during the autumn migration. Thus, parasite transmission by *Culicoides* vectors, for forest-dwelling falconids, may happen mainly after fledging and before the first autumn migration. This seems not to be the case for chicks of the Eurasian Sparrowhawk (*Accipiter nisus*) which showed very high *Leucocytozoon toddi* parasitemias as early as 12–14 d of age (Peirce and Marquiss 1983, Ashford et al. 1991) as well as for chicks of the Northern Goshawk (*Accipiter gentilis*) (Toyne and Ashford 1997). Tella et al. (1999) found that macrohabitat constraints are important in the dynamics of hematozoan transmission. They suggested that the overall low prevalence of blood parasites in Spanish diurnal birds of prey may be due to an overall scarcity of hemoparasite vectors, because Iberian habitats are commonly drier and less-forested than temperate or boreal areas. Sol et al. (2000) have shown that the prevalence of *Haemoproteus columbae*, among near-by populations of the Rock Dove (*Columba livia*), unequivocally paralleled variation in abundance of its main vector, which represents strong support for the hypothesis linking prevalence and vector abundance.

A more comprehensive insect survey and the sampling of adult birds in their breeding colonies would be required to strengthen our conclusions.

RESUMEN.—Se examinaron 42 frotis sanguíneos de pollos volantes de halcón de Eleonora (*Falco eleonora*) de la colonia de las islas Columbretes (NE, España). No se hallaron hemoparásitos en ninguna de las muestras. Se sugiere que la ausencia de vectores apropiados en las islas podría explicar la ausencia de parásitos sanguíneos, aunque se deberá realizar un mayor esfuerzo de muestreo

de vectores así como obtener frotis de aves adultas para confirmar nuestras conclusiones.

[Traducción de los autores]

ACKNOWLEDGMENTS

This study is the contribution No. 4 to the LIFE-NATURE program BA 3200/98/447 "Conservation of island SPAs in the Valencian Region" financed by the Generalitat Valenciana and the E.U. Robert E. Ricklefs provided lab material and allowed us to use the microscope room in the Biology Department of the University of Missouri-St. Louis for this study. Cati and Elena from the Centro de Estudio y Protección del Medio Natural also provided some lab material and helpful advice for blood sampling. G. Urios received financial aid from the Generalitat Valenciana. Santiago Merino reviewed an advanced version of the manuscript. Our special gratitude to all the wardens of the Columbretes Islands.

LITERATURE CITED

- APANIUS, V. 1991. Blood parasitism, immunity and reproduction in American Kestrels (*Falco sparverius*). Ph.D. dissertation, Univ. of Pennsylvania, PA U.S.A.
- AND C.E. KIRKPATRICK. 1988. Preliminary report of *Haemoproteus tinnunculi* infection in a breeding population of American Kestrels (*Falco sparverius*). *J. Wildl. Dis.* 24:150–153.
- ASHFORD, R.W., E.E. GREEN, P.R. HOLMES, AND A.J. LUCAS. 1991. *Leucocytozoon toddi* in British Sparrowhawks *Accipiter nisus*: patterns of infection in nestlings. *J. Nat. Hist.* 25:269–277.
- BLANCO, G., A. GAJON, G. DOVAL, AND F. MARTINEZ. 1998. Absence of blood parasites in Griffon Vultures from Spain. *J. Wildl. Dis.* 34:640–643.
- BENNETT, G.F., R.A. EARLE, H. DU TOIT, AND F.W. HUCHZERMAYER. 1992. A host-parasite catalogue of the haematozoa of the Sub-Saharan birds. *Onderstepoort J. Vet. Res.* 59:1–73.
- BISHOP, M.A. AND G.F. BENNETT. 1992. Host-parasite catalogue of the avian hematozoa. Suppl. 1. Bibliography of the avian blood-inhabiting hematozoa. Suppl. 2. Occas. Pap. Biol. No. 15.
- CASTELLUCCI, S.A., S.B. OLINGER, AND J.R. KLUCSARITS. 1998. Hematology and detection of hemoparasites in the American Kestrel (*Falco sparverius*) during summer nesting period. *J. Pa. Acad. Sci.* 72:29–31.
- FIGUEROLA, J., R. VELARDE, A. BERTOLERO, AND F. CERDA. 1996. Absence of hematozoa in a breeding population of Kentish Plover *Charadrius alexandrinus* in northeast Spain. *J. Ornithol.* 137:523–525.
- AND A. GREEN. 2000. Haematozoan parasites and migratory behaviour in waterfowl. *Evol. Ecol.* 14:143–153.
- GREINER, E.C., G.F. BENNETT, E.M. WHITE, AND R.F. COOMBS. 1975. Distribution of the avian hematozoa of North America. *Can. J. Zool.* 53:1762–1787.
- KIMSEY, R.B. AND P.B. CHANIOTIS. 1984. A light trap for

- bitting Nematocera in moist environments. *Mosq. News* 44:408–412.
- KORPIMÄKI, E., P. TOLONEN, AND G.F. BENNET. 1995. Blood parasites, sexual selection and reproductive success of European Kestrels. *Ecoscience* 2:335–343.
- LITTLE, R.M. AND R.A. EARLÉ. 1994. Lack of avian haematozoa in the Phasianinae of Robben Island. *Ostrich* 65:343–344.
- PEIRCE, M.A., G.F. BENNETT, AND M. BISHOP. 1990. The haemoproteids of the avian order Falconiformes. *J. Nat. Hist.* 24:1091–1100.
- AND M. MARQUISS. 1983. Haematozoa of British birds. VII. Haematozoa of raptors in Scotland with a description of *Haemaproteus nisi* sp. Nov. from the Sparrowhawk (*Accipiter nisus*). *J. Nat. Hist.* 17:813–821.
- PIERSMA, T. 1997. Do global patterns of habitat use and migration strategies co-evolve with relative investments in immunocompetence due to spatial variation in parasite pressure? *Oikos* 80:623–631.
- RISTOW, D. AND M. WINK. 1985. Breeding success and conservation management of Eleonora's Falcon. ICBP Tech. Publ. 5.
- SHELDON, B.C. AND S. VEHLST. 1996. Ecological immunology: costly parasite defenses and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* 11:317–321.
- SOL, D., R. JOVANI, AND J. TORRES. 2000. Geographical variation in blood parasites in feral pigeons: the role of vectors. *Ecography* 23:307–314.
- TELLA, J.L., M. FORERO, A. GAJON, F. HIRALDO, AND J.A. DONAZAR. 1996. Absence of blood-parasitization effects on Lesser Kestrel fitness. *Auk* 113:253–256.
- , G. BLANCO, M. FORERO, A. GAJON, J.A. DONAZAR, AND F. HIRALDO. 1999. Habitat, world geographic range and embryonic development of host explain the prevalence of avian hematozoa at small spatial and phylogenetic scales. *Proc. Nat. Acad. Sci.* 96:1785–1789.
- TOYNE, E.P. AND R.W. ASHFORD. 1997. Blood parasites of nestlings Goshawks. *J. Raptor Res.* 31:81–83.
- WALTER, H. 1979. Eleonora's Falcon. Adaptations to prey and habitat in a social raptor. The Univ. of Chicago Press, Chicago, IL U.S.A.
- WIEHN, J., E. KORPIMÄKI, K.L. BILDSTEIN, AND J. SORJONEN. 1997. Mate choice and reproductive success in the American Kestrel: a role for blood parasites? *Ethology* 103:304–317.
- WINK, M., D. RISTOW, AND C. WINK. 1979. Parasitaemia of adult and juvenile falcons in relation to breeding season and growth. *J. Field Ornithol.* 120:94–97.

Received 10 March 2001; accepted 6 January 2002

J. Raptor Res. 36(2):141–143

© 2002 The Raptor Research Foundation, Inc.

POSSIBLE CHOKING MORTALITIES OF ADULT NORTHERN GOSHAWKS

THOMAS D. BLOXTON¹

Wildlife Science Group, College of Forest Resources, University of Washington, Seattle, WA 98195 U.S.A.

ANDI ROGERS

Arizona Cooperative Fish and Wildlife Research Unit, Biological Sciences East, University of Arizona, Tucson, AZ 85721 U.S.A.

MICHAEL F. INGRALDI AND STEVE ROSENSTOCK

Research Branch, Arizona Game and Fish Department, 2221 West Greenway Road, Phoenix, AZ 85023 U.S.A.

JOHN M. MARZLUFF

Wildlife Science Group, College of Forest Resources, University of Washington, Seattle, WA 98195 U.S.A.

SEAN P. FINN

Boise State University, 1910 University Drive, Boise, ID 83725 U.S.A.

KEY WORDS: Northern Goshawk; *Accipiter gentilis*; choking; mortality; prey.

Choking deaths in wild birds are rarely reported in the ornithological literature. Such incidences have been reported in some easily-observed birds such as Pelecaniformes (Skead 1980, Wilson and Wilson 1985, Septon 1989, Bunkley et al. 1994) and Anseriformes (Septon 1989,

¹ E-mail address: tblox@u.washington.edu